arXiv:1411.7244v1 [math.CA] 26 Nov 2014

ON THE GENERALIZED DIXON INTEGRAL EQUATION

SEMYON YAKUBOVICH*

Department of Mathematics, Faculty of Sciences,
University of Porto, Campo Alegre st., 687, 4169-007 Porto, Portugal

ABSTRACT. A unique analytic solution of the generalized Dixon nonhomogeneous integral equation is derived in $C^1[0,A]$, $A>0$. It is written in terms of the Neumann series, which is expressed as a double series of residues at multiple poles of powers of the gamma-function.

Keywords: *Dixon integral equation, Mellin transform, Gamma-function, Beta- function, Neumann series*

AMS Subject Classifications: 44A15, 45E10

Consider the following nonhomogeneous integral equation

$$f(x)=1+\lambda\int_0^A K(x,y)f(y)dy,\ x\in(0,A),\ \lambda\in\mathbb{C}\setminus\{0\},\ A\geq 1, \tag{1}$$

where

$$K(x,y)=\frac{\partial}{\partial y}P\left(\frac{y}{x}\right)$$

and

$$P(x)=\frac{1}{B(a,a+1)}\int_0^x\frac{t^{a-1}}{(1+t)^{1+2a}}\,dt,\ x\geq 0,\ a>0,$$

where is the Euler beta-function [1], Vol. I. This equation is a generalization of the familiar Dixon integral equation [2], Chapter 11. Let $f\in C^1[0,A]$. Differentiating both sides of (1) with respect to x and then integrating by parts, we find for $x>0$

$$\begin{aligned}f'(x)&=\lambda\frac{d}{dx}\left[P\left(\frac{y}{x}\right)f(y)\,|_0^A-\int_0^A P\left(\frac{y}{x}\right)f'(y)dy\right]\\&=-\frac{\lambda\,x^{-a-1}}{B(a,a+1)}\left(1+\frac{y}{x}\right)^{-1-2a}y^a f(y)\,|_0^A\\&+\frac{\lambda\,x^{-a-1}}{B(a,a+1)}\int_0^A\left(\frac{x}{x+y}\right)^{1+2a}y^a f'(y)dy,\end{aligned} \tag{2}$$

where the differentiation under the integral sign is possible for all $x\geq 0$ due to the absolute and uniform convergence. Indeed, we have

$$x^{-a-1}\int_0^A\left(\frac{x}{x+y}\right)^{1+2a}y^a|f'(y)|dy\leq\max_{x\in[0,A]}|f'(x)|\int_0^{A/x}\frac{y^a}{(1+y)^{1+2a}}dy$$

E-mail: syakubov@fc.up.pt.

$$\leq \max_{x\in[0,A]}|f'(x)|\int_0^\infty \frac{y^a}{(1+y)^{1+2a}}dy<\infty. \tag{3}$$

Moreover, the integrated term vanishes in $y=0$. Therefore we get from (2) the equation

$$f'(x)=-\frac{\lambda f(A)A^a}{B(a,a+1)}\frac{x^a}{(x+A)^{1+2a}}$$

$$+\frac{\lambda}{B(a,a+1)}\int_0^A\left(\frac{x}{y}\right)^a\left(1+\frac{x}{y}\right)^{-1-2a}f'(y)\frac{dy}{y}. \tag{4}$$

Evidently, $f(x)\equiv 0$ does not satisfy the equation (1). Moreover, we will show that $f(x)\equiv$ const is not a solution of (1). Indeed, since (1) can be written in the form

$$f(x)=1+\frac{\lambda x^{a+1}}{B(a,a+1)}\int_0^A\frac{y^{a-1}}{(x+y)^{1+2a}}f(y)dy, \tag{5}$$

assuming that $f(x)=C$ is a solution, we have from (5)

$$C-1=\frac{\lambda C}{B(a,a+1)}\int_0^{A/x}\frac{y^{a-1}}{(1+y)^{1+2a}}dy,\ x>0.$$

Differentiating both sides of the latter equality with respect to x, we find

$$-\frac{\lambda C\,A}{B(a,a+1)x^2}\left(\frac{A}{x}\right)^{a-1}\left(1+\frac{A}{x}\right)^{-1-2a}=0$$

for all $x>0$, which means that $C=0$. But as we saw this is impossible. Further, similar to (3) using the Weierstrass test of the uniform convergence, we verify that the integral in (5) converges absolutely and uniformly for all $x\geq 0$. Hence we find the limit values $f(0)=1$ and $\lim_{x\to+\infty}f(x)=1$.

Let $g(x)=f'(x)H(A-x)$, where $H(x)$ is the Heaviside function. Then applying the Mellin transform [2], [3], [4], Vol. 3

$$g^*(s)=\int_0^\infty g(x)x^{s-1}dx,$$

to both sides of the equation (4), we take into account that the integral in (4) is the convolution for the Mellin transform of $g(x)$ [2] with the function

$$h(x)=\frac{x^a}{(1+x)^{1+2a}},$$

whose Mellin's transform is calculated via the known beta-integral [4], Vol. I

$$\int_0^\infty\frac{x^{a-1}}{(1+x)^{1+2a}}\,dx=B(a,a+1),\ a>0. \tag{6}$$

Thus we obtain $h^*(s)=B(a+s,a+1-s)$. Consequently, after application to (4) of the Mellin transform and the change of the order of integration by Fubini's theorem by virtue of the estimate

$$\int_0^\infty x^{\sigma-1}dx\int_0^\infty\left(\frac{x}{y}\right)^a\left(1+\frac{x}{y}\right)^{-1-2a}|g(y)|\frac{dy}{y}$$

$$=\int_0^\infty\frac{x^{\sigma+a-1}}{(1+x)^{1+2a}}dx\int_0^A y^{\sigma-1}|f'(y)|dy$$

$$\leq\max_{x\in[0,A]}|f'(x)|\int_0^\infty\frac{x^{\sigma+a-1}}{(1+x)^{1+2a}}dx\int_0^A y^{a-1}dy<\infty$$

where $\sigma = \text{Re s}$, $-a < \sigma < a+1$, we come up with the algebraic equality

$$g^*(s)\left[1 - \frac{\lambda\, B(a+s,a+1-s)}{B(a,a+1)}\right]$$
$$= -\lambda\, f(A)A^{s-1}\,\frac{B(a+s,a+1-s)}{B(a,a+1)}, \quad -a < \text{Re s} < a+1. \tag{7}$$

Therefore, if

$$\left|\frac{\lambda\, B(a+s,a+1-s)}{B(a,a+1)}\right| \le \frac{|\lambda|\, B(a+\sigma,a+1-\sigma)}{B(a,a+1)} < 1$$

i.e.

$$|\lambda| < \frac{B(a,a+1)}{B(a+\sigma,a+1-\sigma)} \tag{8}$$

equality (7) becomes

$$g^*(s) = -\lambda\, f(A)A^{s-1}\,\frac{B(a+s,a+1-s)}{B(a,a+1) - \lambda B(a+s,a+1-s)}. \tag{9}$$

Now, considering $-a < \sigma < a+1$, we use the inverse Mellin transform [2] to provide the unique solution of the equation (4) in the form

$$f'(x) = -\frac{\lambda\, f(A)}{2\pi A i}\int_{\sigma-i\infty}^{\sigma+i\infty}\frac{B(a+s,a+1-s)}{B(a,a+1) - \lambda B(a+s,a+1-s)}\left(\frac{x}{A}\right)^{-s} ds. \tag{10}$$

Let $\sigma \neq 1$. Then if $-a < \sigma < 1$, we integrate both sides of (10) from 0 to x, and taking into account account the value $f(0) = 1$, we come up with the unique soliton of the original integral equation (1), namely

$$f(x) = 1 - \frac{\lambda\, f(A)}{2\pi i}\int_{\sigma-i\infty}^{\sigma+i\infty}\frac{B(a+s,a+1-s)}{B(a,a+1) - \lambda B(a+s,a+1-s)}\left(\frac{x}{A}\right)^{1-s}\frac{ds}{1-s}. \tag{11}$$

Our final goal is to write the solution (11) in a different form, using the corresponding Neumann series and calculating the Mellin-Barnes integral by the residue theorem at multiple poles of powers of the gamma-function [1], Vol. 1, [4], Vol. 3. To do this, we will employ the series representation of the gamma-function $\Gamma(z)$ near the poles $z = -m$, $m \in \mathbb{N}_0$ (see, for instance, in [5])

$$\Gamma(z) = \frac{(-1)^m}{z+m}\sum_{k=0}^{\infty} c_{k,m}(z+m)^k, \tag{12}$$

where

$$c_{k,m} = \sum_{v=0}^{k}\frac{(-1)^{(v+k)/2-1}(2^{k-v}-2)B_{k-v}\pi^{k-v}}{v!(k-v)!}d_{v,m}, \tag{13}$$
$$d_{v,m} = \lim_{w\to m+1}\frac{d^v}{dw^v}\left[\frac{1}{\Gamma(w)}\right]$$

and B_μ are the Bernoulli numbers [1], Vol. I. We begin, writing (11) in the form

$$f(x) = 1 - \frac{f(A)}{2\pi i}\int_{\sigma-i\infty}^{\sigma+i\infty}\sum_{n=1}^{\infty}\left(\frac{\lambda\, B(a+s,a+1-s)}{B(a,a+1)}\right)^n\left(\frac{x}{A}\right)^{1-s}\frac{ds}{1-s}. \tag{14}$$

The change of the order of integration and summation is allowed owing to the absolute convergence which, in turn, can be verified by virtue of the Stirling asymptotic formula for the gamma-function [1], Vol. I , condition (7) and the estimate

$$\int_{\sigma-i\infty}^{\sigma+i\infty}\sum_{n=1}^{\infty}\left|\left(\frac{\lambda\, B(a+s,a+1-s)}{B(a,a+1)}\right)^n\left(\frac{x}{A}\right)^{1-s}\frac{ds}{1-s}\right|$$

$$\leq \left(\frac{x}{A}\right)^{1-\sigma}\sum_{n=0}^{\infty}\left(\frac{|\lambda|\,B(a+\sigma,a+1-\sigma)}{B(a,a+1)}\right)^{n}\int_{\sigma-i\infty}^{\sigma+i\infty}|B(a+s,a+1-s)|\frac{|ds|}{|1-s|}<\infty,$$

where $\sigma\in(-a,1)$. Therefore we write solution (14) as

$$f(x)=1-f(A)\sum_{n=1}^{\infty}\left(\frac{\lambda}{\Gamma(a)\Gamma(a+1)}\right)^{n}\frac{1}{2\pi i}\int_{\sigma-i\infty}^{\sigma+i\infty}[\Gamma(a+s)\Gamma(a+1-s)]^{n}\left(\frac{x}{A}\right)^{1-s}\frac{ds}{1-s}. \tag{15}$$

Meanwhile, the integral in (15) can be calculated by a special version of the residue theorem, namely, the Slater theorem (see [4], Vol. 3) as the series of residues of the integrand at multiple left-hand poles $s=-a-m,\ m\in\mathbb{N}_0$ of the powers $[\Gamma(a+s)]^n,\ n\in\mathbb{N}$. We have

$$\frac{1}{2\pi i}\int_{\sigma-i\infty}^{\sigma+i\infty}[\Gamma(a+s)\Gamma(a+1-s)]^{n}\left(\frac{x}{A}\right)^{1-s}\frac{ds}{1-s}$$
$$=\sum_{m=0}^{\infty}\mathrm{Res}_{s=-a-m}\left[[\Gamma(a+s)\Gamma(a+1-s)]^{n}\left(\frac{x}{A}\right)^{1-s}\frac{1}{1-s}\right]. \tag{16}$$

However,

$$\mathrm{Res}_{s=-a-m}\left[[\Gamma(a+s)\Gamma(a+1-s)]^{n}\left(\frac{x}{A}\right)^{1-s}\frac{1}{1-s}\right]$$
$$=\frac{1}{(n-1)!}\lim_{s\to-a-m}\frac{d^{n-1}}{ds^{n-1}}\left[[(s+a+m)\Gamma(a+s)]^{n}[\Gamma(a+1-s)]^{n}\left(\frac{x}{A}\right)^{1-s}\frac{1}{1-s}\right]$$
$$=\frac{1}{(n-1)!}\lim_{s\to-a-m}\sum_{r=0}^{n-1}\binom{n-1}{r}\left[[(s+a+m)\Gamma(a+s)]^{n}\right]^{(r)}\left[[\Gamma(a+1-s)]^{n}\left(\frac{x}{A}\right)^{1-s}\frac{1}{1-s}\right]^{(n-1-r)}. \tag{17}$$

In the meantime,

$$\left[[\Gamma(a+1-s)]^{n}\left(\frac{x}{A}\right)^{1-s}\frac{1}{1-s}\right]^{(n-1-r)}$$
$$=\sum_{\nu=0}^{n-1-r}\binom{n-1-r}{\nu}([\Gamma(a+1-s)]^{n})^{(\nu)}\left(\left(\frac{x}{A}\right)^{1-s}\frac{1}{1-s}\right)^{(n-1-r-\nu)}$$
$$=\left(\frac{x}{A}\right)^{1-s}(n-1-r)!\sum_{\nu=0}^{n-1-r}\frac{(-1)^{n-1-r-\nu}}{\nu!}([\Gamma(a+1-s)]^{n})^{(\nu)}$$
$$\times\sum_{k=0}^{n-1-r-\nu}\frac{(-1)^{k}(\log(x/A))^{n-1-r-\nu-k}}{(n-1-r-\nu-k)!(1-s)^{k+1}}. \tag{18}$$

Meanwhile, in order to calculate higher -order derivatives of powers $[(s+a+m)\Gamma(a+s)]^n$, $[\Gamma(a+1-s)]^n$ we will appeal to the familiar Faá di Bruno formula [6]. Thus we obtain

$$\left[[(s+a+m)\Gamma(a+s)]^{n}\right]^{(r)}=\sum\frac{r!\ n!\ ((s+a+m)\Gamma(a+s))^{n-l}}{(n-l)!b_1!b_2!\dots b_r!}\left(\frac{((s+a+m)\Gamma(a+s))^{(1)}}{1!}\right)^{b_1}\dots$$
$$\times\left(\frac{((s+a+m)\Gamma(a+s))^{(r)})}{r!}\right)^{b_r}, \tag{19}$$

where the sum (19) is over all different solutions in nonnegative integers $b_1,b_2,\dots,b_r$ of $b_1+2b_2+\dots+rb_r=r$ and $l=b_1+b_2+\dots+b_r$ and

$$\left[[\Gamma(a+1-s)]^{n}\right]^{(\nu)}=\sum\frac{\nu!\ n!\ (\Gamma(a+1-s))^{n-l}}{(n-q)!\beta_1!\beta_2!\dots\beta_\nu!}\left(\frac{\Gamma^{(1)}(a+1-s)}{1!}\right)^{\beta_1}\dots\left(\frac{\Gamma^{(\nu)}(a+1-s)}{\nu!}\right)^{b_\nu}, \tag{20}$$

where the sum (20) is over all different solutions in nonnegative integers $\beta_1,\beta_2,\dots,\beta_\nu$ of $\beta_1+2\beta_2+\dots+r\beta_\nu=\nu$ and $q=\beta_1+\beta_2+\dots+\beta_\nu$. Hence, recalling the expansion (12) of the gamma-function near the poles, we find, correspondingly,

$$\lim_{s\to -a-m}[[(s+a+m)\Gamma(a+s)]^n]^{(r)}=\sum \frac{r!\ n!\ (-1)^{mn}\ c_{1,m}^{b_1}\dots c_{r,m}^{b_r}}{(n-l)!(m!)^{n-l}b_1!b_2!\dots b_r!}, \tag{21}$$

$$\lim_{s\to -a-m}[[\Gamma(a+1-s)]^n]^{(\nu)}=\sum \frac{\nu!\ n!\ (\Gamma(2a+1+m))^{n-q}}{(n-q)!\beta_1!\beta_2!\dots\beta_\nu!}\left(\frac{\Gamma^{(1)}(2a+1+m)}{1!}\right)^{\beta_1}\dots\left(\frac{\Gamma^{(\nu)}(2a+1+m)}{\nu!}\right)^{\beta_\nu}. \tag{22}$$

Consequently, combining with (17), (18), we obtain values of the residues

$$\mathrm{Res}_{s=-a-m}\left[[\Gamma(a+s)\Gamma(a+1-s)]^n\left(\frac{x}{A}\right)^{1-s}\frac{1}{1-s}\right]=(n!)^2\left(\frac{x}{A}\right)^{a+1+m}\sum_{r=0}^{n-1}\sum_{\nu=0}^{n-1-r}(-1)^{n-1-r-\nu}$$

$$\times\sum_{k=0}^{n-1-r-\nu}\frac{(-1)^k(\log(x/A))^{n-1-r-\nu-k}}{(n-1-r-\nu-k)!(a+1+m)^{k+1}}\sum_b\sum_\beta\frac{(-1)^{mn}\ c_{1,m}^{b_1}\dots c_{r,m}^{b_r}}{(n-l)!(m!)^{n-l}b_1!b_2!\dots b_r!}\frac{\Gamma^{n-q}(2a+1+m)}{(n-q)!\beta_1!\beta_2!\dots\beta_\nu!}$$

$$\times\left(\frac{\Gamma^{(1)}(2a+1+m)}{1!}\right)^{\beta_1}\dots\left(\frac{\Gamma^{(\nu)}(2a+1+m)}{\nu!}\right)^{\beta_\nu}. \tag{23}$$

Thus the value of the integral (16) is equal to

$$\frac{1}{2\pi i}\int_{\sigma-i\infty}^{\sigma+i\infty}[\Gamma(a+s)\Gamma(a+1-s)]^n\left(\frac{x}{A}\right)^{1-s}\frac{ds}{1-s}$$

$$=(n!)^2\sum_{m=0}^{\infty}\sum_{r=0}^{n-1}\sum_{\nu=0}^{n-1-r}(-1)^{n-1-r-\nu}$$

$$\times\left(\frac{x}{A}\right)^{a+1+m}\sum_{k=0}^{n-1-r-\nu}\frac{(-1)^k\ (\log(x/A))^{n-1-r-\nu-k}}{(n-1-r-\nu-k)!(a+1+m)^{k+1}}$$

$$\times\sum_b\sum_\beta\frac{(-1)^{mn}\ c_{1,m}^{b_1}\dots c_{r,m}^{b_r}}{(n-l)!(m!)^{n-l}b_1!b_2!\dots b_r!}\frac{\Gamma^{n-q}(2a+1+m)}{(n-q)!\beta_1!\beta_2!\dots\beta_\nu!}$$

$$\times\left(\frac{\Gamma^{(1)}(2a+1+m)}{1!}\right)^{\beta_1}\dots\left(\frac{\Gamma^{(\nu)}(2a+1+m)}{\nu!}\right)^{\beta_\nu} \tag{24}$$

and therefore our unique solution of the equation (1) takes the form for $x\in[0,A]$

$$f(x)=1-f(A)\left(\frac{x}{A}\right)^{a+1}\sum_{n=1}^{\infty}\sum_{m=0}^{\infty}\left(\frac{(-1)^m\lambda}{\Gamma(a)\Gamma(a+1)}\right)^n\left(\frac{x}{A}\right)^m\sum_{r=0}^{n-1}\sum_{\nu=0}^{n-1-r}(-1)^{n-1-r-\nu}$$

$$\times\sum_{k=0}^{n-1-r-\nu}\frac{(-1)^k\ (\log(x/A))^{n-1-r-\nu-k}}{(n-1-r-\nu-k)!(a+1+m)^{k+1}}\sum_b\sum_\beta\binom{n}{l}\binom{n}{q}\frac{l!\ q!\ c_{1,m}^{b_1}\dots c_{r,m}^{b_r}\ \Gamma^{n-q}(2a+1+m)}{(m!)^{n-l}\ b_1!b_2!\dots\ b_r!\ \beta_1!\beta_2!\dots\beta_\nu!}$$

$$\times\left(\frac{\Gamma^{(1)}(2a+1+m)}{1!}\right)^{\beta_1}\dots\left(\frac{\Gamma^{(\nu)}(2a+1+m)}{\nu!}\right)^{\beta_\nu}. \tag{25}$$

Further, suppose that $1<\sigma<a+1$. Then recalling (10), we integrate from x to A to get the equality

$$f(x)-f(A)=\frac{\lambda f(A)}{2\pi i}\int_{\sigma-i\infty}^{\sigma+i\infty}\frac{B(a+s,a+1-s)}{B(a,a+1)-\lambda B(a+s,a+1-s)}\left[1-\left(\frac{x}{A}\right)^{1-s}\right]\frac{ds}{1-s}. \tag{26}$$

Hence, in particular, passing to the limit in (26) when $x\to+\infty$, we remind the value $\lim_{x\to+\infty}f(x)=1$ to derive the equality

$$f(A)\left[1+\frac{\lambda}{2\pi i}\int_{\sigma-i\infty}^{\sigma+i\infty}\frac{B(a+s,a+1-s)}{B(a,a+1)-\lambda B(a+s,a+1-s)}\frac{ds}{1-s}\right]=1. \tag{27}$$

Now, following the same scheme as above, we extend our solution for $x\geq A$. Indeed, making a simple substitution, we have (see (17), (18))

$$\frac{1}{2\pi i}\int_{\sigma-i\infty}^{\sigma+i\infty}[\Gamma(a+s)\Gamma(a+1-s)]^n\left[1-\left(\frac{x}{A}\right)^{1-s}\right]\frac{ds}{1-s}$$

$$=\frac{1}{2\pi i}\int_{1-\sigma-i\infty}^{1-\sigma+i\infty}[\Gamma(a+s)\Gamma(a+1-s)]^n\left[1-\left(\frac{x}{A}\right)^{s}\right]\frac{ds}{s}$$

$$=\sum_{m=0}^{\infty}\mathrm{Res}_{s=-a-m}\left[[\Gamma(a+s)\Gamma(a+1-s)]^n\left[1-\left(\frac{x}{A}\right)^{s}\right]\frac{1}{s}\right]$$

$$=-\sum_{m=0}^{\infty}(-1)^{mn}\sum_{r=0}^{n-1}\sum_{\nu=0}^{n-1-r}\left[\frac{1}{(a+m)^{n-r-\nu}}-\left(\frac{x}{A}\right)^{-a-m}\sum_{k=0}^{n-1-r-\nu}\frac{(\log(x/A))^{n-1-r-\nu-k}}{(n-1-r-\nu-k)!(a+m)^{k+1}}\right]$$

$$\times\sum_{b}\sum_{\beta}\binom{n}{l}\binom{n}{q}\frac{l!\,q!\,c_{1,m}^{b_1}\dots c_{r,m}^{b_r}\,\Gamma^{n-q}(2a+1+m)}{(m!)^{n-l}\,b_1!b_2!\dots\,b_r!\,\beta_1!\beta_2!\dots\beta_\nu!}\left(\frac{\Gamma^{(1)}(2a+1+m)}{1!}\right)^{\beta_1}\dots\left(\frac{\Gamma^{(\nu)}(2a+1+m)}{\nu!}\right)^{\beta_\nu}$$

and (26) becomes $(x\geq A)$

$$f(x)=f(A)\left[1-\sum_{n=1}^{\infty}\sum_{m=0}^{\infty}\left(\frac{\lambda(-1)^m}{\Gamma(a)\Gamma(a+1)}\right)^n\sum_{r=0}^{n-1}\sum_{\nu=0}^{n-1-r}\left[\frac{1}{(a+m)^{n-r-\nu}}\right.\right.$$

$$\left.\left.-\left(\frac{x}{A}\right)^{-a-m}\sum_{k=0}^{n-1-r-\nu}\frac{(\log(x/A))^{n-1-r-\nu-k}}{(n-1-r-\nu-k)!(a+m)^{k+1}}\right]\right.$$

$$\left.\times\sum_{b}\sum_{\beta}\binom{n}{l}\binom{n}{q}\frac{l!\,q!\,c_{1,m}^{b_1}\dots c_{r,m}^{b_r}\,\Gamma^{n-q}(2a+1+m)}{(m!)^{n-l}\,b_1!b_2!\dots\,b_r!\,\beta_1!\beta_2!\dots\beta_\nu!}\left(\frac{\Gamma^{(1)}(2a+1+m)}{1!}\right)^{\beta_1}\dots\left(\frac{\Gamma^{(\nu)}(2a+1+m)}{\nu!}\right)^{\beta_\nu}\right].$$

We summarize our results by the following

Theorem. *Let $f\in C^1[0,A]$, $A\geq 1$, $\lambda\in\mathbb{C}\setminus\{0\}$, satisfying the condition*

$$|\lambda|<\frac{B(a,a+1)}{B(a+\sigma,a+1-\sigma)},\ a>0,$$

where $\sigma\in(-a,1+a)\setminus\{1\}$. Then the integral equation

$$f(x)=1+\lambda\int_0^A K(x,y)f(y)dy,$$

where

$$K(x,y)=\frac{\partial}{\partial y}P\left(\frac{y}{x}\right)$$

and

$$P(x)=\frac{1}{B(a,a+1)}\int_0^x \frac{t^{a-1}}{(1+t)^{1+2a}}\,dt$$

has the unique solution represented by the formula

$$f(x)=1-f(A)\left(\frac{x}{A}\right)^{a+1}\sum_{n=1}^{\infty}\sum_{m=0}^{\infty}\left(\frac{(-1)^m\lambda}{\Gamma(a)\Gamma(a+1)}\right)^n\left(\frac{x}{A}\right)^m\sum_{r=0}^{n-1}\sum_{\nu=0}^{n-1-r}(-1)^{n-1-r-\nu}$$
$$\times\sum_{k=0}^{n-1-r-\nu}\frac{(-1)^k\,(\log(x/A))^{n-1-r-\nu-k}}{(n-1-r-\nu-k)!(a+1+m)^{k+1}}\sum_b\sum_\beta\binom{n}{l}\binom{n}{q}\frac{l!\,q!\,c_{1,m}^{b_1}\dots c_{r,m}^{b_r}\,\Gamma^{n-q}(2a+1+m)}{(m!)^{n-l}\,b_1!b_2!\dots\,b_r!\,\beta_1!\beta_2!\dots\beta_\nu!}$$
$$\times\left(\frac{\Gamma^{(1)}(2a+1+m)}{1!}\right)^{\beta_1}\dots\left(\frac{\Gamma^{(\nu)}(2a+1+m)}{\nu!}\right)^{\beta_\nu},$$

where summations over b and β are explained in (20) *and $c_{j,m}$ are defined by* (13)*. Moreover, it can be extended analytically for $x\geq A$ by the equality*

$$f(x)=f(A)\left[1-\sum_{n=1}^{\infty}\sum_{m=0}^{\infty}\left(\frac{\lambda(-1)^m}{\Gamma(a)\Gamma(a+1)}\right)^n\sum_{r=0}^{n-1}\sum_{\nu=0}^{n-1-r}\left[\frac{1}{(a+m)^{n-r-\nu}}\right.\right.$$
$$\left.\left.-\left(\frac{x}{A}\right)^{-a-m}\sum_{k=0}^{n-1-r-\nu}\frac{(\log(x/A))^{n-1-r-\nu-k}}{(n-1-r-\nu-k)!(a+m)^{k+1}}\right]\right.$$
$$\left.\times\sum_b\sum_\beta\binom{n}{l}\binom{n}{q}\frac{l!\,q!\,c_{1,m}^{b_1}\dots c_{r,m}^{b_r}\,\Gamma^{n-q}(2a+1+m)}{(m!)^{n-l}\,b_1!b_2!\dots\,b_r!\,\beta_1!\beta_2!\dots\beta_\nu!}\left(\frac{\Gamma^{(1)}(2a+1+m)}{1!}\right)^{\beta_1}\dots\left(\frac{\Gamma^{(\nu)}(2a+1+m)}{\nu!}\right)^{\beta_\nu}\right].$$

Acknowledgments

I like to thank A. Polunchenko and I. Petrov for bringing this integral equation and its important applications in Statistics to my attention.

References

1. A. Erdélyi, W. Magnus, F. Oberhettinger and F.G. Tricomi, *Higher Transcendental Functions*, Vols. I and II, McGraw-Hill, New York, London and Toronto (1953).
2. E.C. Titchmarsh, *An Introduction to the Theory of Fourier Integrals*, Clarendon Press, Oxford (1937).
3. S. Yakubovich and Yu. Luchko, *The Hypergeometric Approach to Integral Transforms and Convolutions*. Mathematics and its Applications, 287. Kluwer Academic Publishers Group, Dordrecht (1994).
4. A.P. Prudnikov, Yu. A. Brychkov and O. I. Marichev, *Integrals and Series: Vol. 1: Elementary Functions*, Gordon and Breach, New York (1986); *Integrals and Series: Vol. 2: Special Functions*, Gordon and Breach, New York (1986); *Vol. 3: More Special Functions*, Gordon and Breach, New York (1990).
5. E.W. Weisstein, "Gamma Function." From MathWorld–A Wolfram Web Resource. http://functions.wolfram.com/GammaBetaErf/Gamma/
6. W.P. Johnson, The curious history of Faá di Bruno's formula, *The Amer. Math.Monthly* , **109** (2002), N 3, 217- 234.